



RESEARCH ARTICLE

The Therapy Process Questionnaire - Factor analysis and psychometric properties of a multidimensional self-rating scale for high-frequency monitoring of psychotherapeutic processes

Günter Schiepek^{1,2,3} | Barbara Stöger-Schmidinger² | Helmut Kronberger² |
Wolfgang Aichhorn^{1,2} | Leonhard Kratzer⁴ | Peter Heinz⁴ | Kathrin Viol^{1,2}  |
Anna Lichtwarck-Aschoff⁵  | Helmut Schöller^{1,2}

¹Institute of Synergetics and Psychotherapy Research, Paracelsus Medical University, Salzburg, Austria

²University Hospital of Psychiatry, Psychotherapy, and Psychosomatics, Paracelsus Medical University, Salzburg, Austria

³Department of Psychology, Ludwig Maximilian University Munich, Munich, Germany

⁴Department of Psychotraumatology, Clinic St. Irmingard, Prien am Chiemsee, Germany

⁵Developmental Psychopathology, Behavioural Science Institute, Radboud University, Nijmegen, The Netherlands

Correspondence

Günter Schiepek, Institute of Synergetics and Psychotherapy Research, Paracelsus Medical University, Salzburg, Austria.
Email: guenter.schiepek@ccsys.de

Abstract

Many outcome measures and session-related questionnaires in psychotherapy are designed for weekly or biweekly administration. Yet, today, technical developments allow for higher frequency assessments to monitor human change dynamics more closely by daily assessments. For this purpose, the Therapy Process Questionnaire (TPQ) was developed, with a specific focus on inpatient psychotherapy. In this article, we present an explorative and confirmative factor analysis of the TPQ on the basis of the time series data of 150 patients collected during their hospital stay (mean time series length: 69.1 measurement points). A seven-factor solution was identified, which explains 68.7% of variance and associates 43 items onto the factors, which are “well-being and positive emotions,” “relationship with fellow patients,” “therapeutic relationship and clinical setting,” “emotional and problem intensity,” “insight/confidence/therapeutic progress,” “motivation for change,” and “mindfulness/self-care.” The internal consistency (Cronbach's α), the inter-item correlations of the subscales, and the discriminative power of the items are excellent. The TPQ can be applied in practice and research for creating time series with equidistant measurement points and time series lengths, which are appropriate for the application of nonlinear analysis methods. Especially in clinical practice, it is important to identify precursors of phase transitions, changing synchronization patterns, and critical or instable periods of a process, which now is possible by internet- or app-based applications of this multidimensional questionnaire.

KEYWORDS

factor analysis, nonlinear dynamics, process monitoring, psychotherapy feedback, Therapy Process Questionnaire

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors *Clinical Psychology & Psychotherapy* Published by John Wiley & Sons Ltd

1 | INTRODUCTION

Outcome monitoring and feedback on therapeutic progress has become popular and has been adopted by many mental health providers all over the world (e.g., Delgado et al., 2017; Gibbons et al., 2015; Schiepek et al., 2016; Wampold, 2015). Lambert (2007, 2010) describes it as an important feature of good clinical practice and asks for an integration of monitoring procedures into routines of mental health care. Studies and meta-analyses report on effects of psychotherapy feedback in outcome optimization ("routine outcome monitoring") at different degrees (mostly small to moderate effect sizes), particularly in patients at risk for deterioration (Brattland et al., 2018; de Jong, van Sluis, Nugter, Heiser, & Spinhoven, 2012; de Jong, Segaar, Ingenhoven, van Busschbach, & Timman, 2018; de Jong, Barkham, Wolpert, & the INSPIRE Research Network, in prep.; Gibbons et al., 2015; Kendrick et al., 2016; Knaup, Kösters, Schöfer, Becker, & Puschner, 2009; Lambert, 2017; Lambert, Whipple, & Kleinstäuber, 2018; Shimokawa, Lambert, & Smart, 2010). Although there is a wide agreement on the usefulness of process and outcome monitoring (Wampold, 2015), there are different ways of applying it to clinical practice. The question of *how* it should be implemented and realized has been answered from different viewpoints.

Diverging perspectives have been previously offered to address the question if human change processes should be conceptualized as linear and input-dependent or as nonlinear, complex, and dynamic. Within the linear approach, treatment courses seem to be more or less continuous, characterized by straight lines or moderately damped trajectories (dose-effect curves), and predictable by standard tracks using the reference of patients with the same diagnosis or the same symptom severity at the beginning (Finch, Lambert, & Schaalje, 2001). In consequence, measures at low frequencies (session by session, weekly, or bi-weekly) seem to be sufficient. Within the nonlinear dynamic systems paradigm (Haken & Schiepek, 2010; Gelo & Salvatore, 2016; Schiepek, Heinzl, Karch, Plöderl, & Strunk, 2016; Schiepek & Pincus, under review), trajectories of change may have complex shapes, are usually characterized by discontinuous pattern transitions, which become evident as sudden gains or losses (Hayes, Laurenceau, Feldman, Strauss, & Cardaciotto, 2007; Lutz et al., 2013; Stiles et al., 2003) or rupture-repair sequences in the working alliance (Gumz, Brähler, Geyer, & Erices, 2012; Stiles et al., 2004), demonstrate critical instabilities and other precursors of discontinuous transitions (Schiepek, Tominschek, & Heinzl, 2014), and are chaotic, that is, not predictable in the long run (Schiepek et al., 2017). In consequence, this creates specific demands on sampling rates and time series length.

Most feedback procedures in psychotherapy are working within the linear low-frequency approach and focus on treatment outcomes. Widely used measures are the Outcome Questionnaire (Lambert et al., 2004), the World Health Organization Wellbeing Index (Bech, Gudex, & Johansen, 1996), the Mental Health subscales of the Medical Outcomes Questionnaire (Ware, Snow, Kosinski, & Gandek, 1993), the Health of the Nation Outcome Scales (Wing et al., 1998), or the Depression Anxiety Stress Scales (Lovibond & Lovibond, 1995;

Key Practitioner Message

- The Therapy Process Questionnaire (TPQ) can be used for daily measurements, which produce continuous time series of psychotherapeutic processes. The questionnaire is created for the administration by electronic devices (app- or internet-based systems).
- This kind of high-frequency assessment allows for the identification of pattern transitions and their precursors (e.g., critical instabilities) as well as other nonlinear features of change dynamics.
- The multidimensional assessment of different mechanisms of change catalyses the personal development of the patients and delivers important information for feedback sessions and the management of the therapeutic process.

Nilges & Essau, 2015). Besides the outcome, process-mediating aspects and known therapeutic mechanisms and mediators also should be respected, for example, client factors (resources, motivation to change, or engagement), working alliance, emotions, self-relatedness, expectancies, self-esteem, or self-efficacy (Duncan, Miller, Wampold, & Hubble, 2010; Norcross & Lambert, 2011; Schiepek, Aichhorn, et al., 2016). There is a diversity of questionnaires focusing on such ingredients of the therapeutic work. Usually, these questionnaires are designed as post-session reports, for example, the Therapy Session Report (Orlinsky & Howard, 1967), the Vanderbilt Psychotherapy Process Scales (Smith, Hilsenroth, Baity, & Knowles, 2003), the Session Evaluation Questionnaire (Stiles, 1980; Stiles et al., 1994), the Session Impact Scale (Elliott & Wexler, 1994), or the Berne Post-Session Report (Flückiger, Regli, Zwahlen, Hostettler, & Caspar, 2010) and its advanced development, the Scale for the Multiperspective Assessment of General Change Mechanisms in Psychotherapy (SACiP; Mander et al., 2013). The multidimensional Questionnaire for the Evaluation of Psychotherapeutic Processes (Lutz & Böhnke, 2008) is applied at some few time points during the process with the ratings referring to a period of 1 week. Other session-related scales focus uniquely on the therapeutic alliance, such as the Working Alliance Inventory (Horvath & Greenberg, 1986), the Helping Alliance Questionnaire (Alexander & Luborsky, 1986), the California Psychotherapy Alliance Scales (Gaston, 1991), or the Combined Alliance Scale (Hatcher & Shannon, 2005) and its short version, the Scale of the Therapeutic Alliance-Revised (STA-R; Brockmann et al., 2011). A measure that was introduced by Mander et al. (2014) combines the items of the SACiP and the STA-R to a further instrument, the Individual Therapy Process Questionnaire. There are also scales that focus on specific constructs or specific types of interventions, such as the Applied Mindfulness Process Scale (Li, Black, & Garland, 2015).

Usually, outcome measures do not cover mechanisms of change and usually grasp longer periods of time (e.g., one, two, or more

weeks). Questionnaires on secondary outcomes (e.g., on the quality of life or interpersonal relations) also focus on specific constructs and grasp longer periods of time. Other process questionnaires are explicitly focused on the experiences of a patient within sessions (usually all alliance questionnaires) or refer to inter-session periods or are linked to specific interventions. In opposition to all possible alternatives, the unique feature of the Therapy Process Questionnaire (TPQ) is its aptitude for high-frequency (daily) monitoring and its capacity to capture different change-related constructs.

Compared with the sampling rate of sessions, which usually take place at irregular time distances, daily measurements appear to be an achievable alternative. Days are more frequent and also equidistant and by this fulfil the criteria of (a) a complete recording of therapies (not only as an irregular event sampling), (b) frequent and (c) continuous measurements, and (d) considering practicalities of data collection. A further point is a therapeutic one: Daily self-assessments—especially when combined with electronic diaries—activate the reasoning on one's own thoughts, feelings, and behaviour (mindfulness and metacognition) and by this support self-regulation and auto-catalyse developmental processes. This is a mechanism of change that is reported by many patients and therapists and is also documented in case studies (e.g., Kratzer et al., 2018; Schiepek, Eckert, Aas, Wallot, & Wallot, 2015; Schiepek, Stöger-Schmidinger, Aichhorn, Schöller, & Aas, 2016; Schiepek, Aichhorn, Schöller, & Kronberger, 2018) and in a controlled study on the effects of process monitoring on chronically addicted inpatients (Patzig & Schiepek, 2015). More systematic studies are needed on the psychological effects of continuous self-reports and on the synergistic effects between monitoring other mechanisms of change.


Figure 1 illustrates how the dynamics of a time series (daily ratings of self-esteem from a patient with borderline personality disorder) is distorted and the information on the dynamic pattern is lost if measurement points are successively omitted. The rapid cycling of self-esteem characterizing the first weeks of a treatment vanishes if ratings are only made on every second (Figure 1b) or fourth day (Figure 1c), weekly (Figure 1d), or at mixed weekly and fortnightly intervals, the most common periodicity of therapy sessions (Figure 1e,f). Corresponding to the loss of information, the dynamics of the presented time series appear more and more linear with the shape of the curve

depending on the chosen measurement points. Following the Nyquist theorem, sampling rates have to be at least twice as high as the eigendynamics of the systems under investigation (Shannon, 1949).

The nonlinear high-frequency approach to psychotherapy takes seriously what in many theoretical and empirical writings was highlighted: The dynamic and non-ergodic nature of human cognitions, emotions, and behaviour needs a switch from linear group statistics to a paradigm of understanding and measuring complex systems by using nonstationary time series analysis and even intra-individual methods, combining idiographic with nomothetic science (Haken & Schiepek, 2010; Molenaar, 2004, 2007; Molenaar, Sinclair, Rovine, Ram, & Corneal, 2009; Piccirillo, Beck, & Rodebaugh, 2019; Schiepek et al., 2015; Schiepek, Heinzl, et al., 2016; Schiepek, Stöger-Schmidinger, et al., 2016; Tzur-Bitan, Meiran, & Shahar, 2010; Tzur-Bitan, Meiran, Steinberg, & Shahar, 2012). Chaotic dynamics produced by the functioning of nonlinear complex systems indeed are not predictable but realize specific dynamic features, which are important for understanding human change. One is the sensitive dependency of the dynamics on minimal input, which creates adaptability and flexibility of the systems to the environment (e.g., therapeutic interventions), another one is the realization of attractors (more or less stable patterns of the dynamics), and a third one is transitions between attractors depending on the control parameters of a system (Strunk & Schiepek, 2006). Within this context, clusters of outcomes may be predicted by any characteristics of psychotherapy, but the concrete trajectories of change will be unpredictable by principal reasons. Instead of including more and more predictors in linear regression models, the consequence will be to create short-term predictions (early warning systems) of critical events or changing dynamics in practice and to create nonlinear system models of involved variables in theory. A sampling rate of daily measures has proven to allow for insight in nonlinear patterns and its transitions (see Figure 1).

About two decades ago, we introduced the TPQ (Haken & Schiepek, 2010; Nischk, Grothe, & Schiepek, 2000) as a method for day-by-day assessments of psychotherapeutic processes. Since about 2 years, an optimized version of the TPQ is available for the routine assessment of psychotherapeutic change dynamics. The optimization was based on many years of clinical experience in inpatient settings.

FIGURE 1 Distortion of the dynamics of a time series by omitting measurement points. The time series represents the self-esteem of a patient with borderline personality disorder during her hospital stay. (a) The original time series with daily measures (opaque in b–f). (b) Every second day is omitted as missing day. Fluctuations of the first weeks of the time series vanish, if ratings are only made on every fourth day (c) or weekly with some variation around an exact 7-day rhythm (d). A major loss of information and possible source of therapeutic misjudgment occurs with the common practice of occasional weekly and fortnightly measurement intervals (e,f).



Despite of the 47 items of the questionnaire, daily ratings never caused big problems. The compliance rates were amazingly high (up to 80%; Schiepek, Aichhorn, et al., 2016), which is due to the facts that (a) the self-reports could be done by electronic devices via the Synergetic Navigation System (SNS), (b) the monitoring was defined as part of the hospital and treatment routine, and—perhaps the most important—(c) the feedbacks on the resulting data (visualized as graphs and diagrams) and analysis results were integrated into the psychotherapeutic process. Patients and therapists refer on these results for planning and evaluating the process (continuous cooperative process control) and would not miss this kind of treatment support. Here, we present a factor analysis and statistical item characteristics of this questionnaire.

2 | METHODS AND MATERIALS

2.1 | Subjects

The patients of this study were treated at two psychotherapy centres, the Department of In-patient Psychotherapy at the University

Hospital of Psychiatry, Psychotherapy and Psychosomatics (Paracelsus Medical University Salzburg, Austria) and the Department of Psychotraumatology at the Clinic St. Irmingard (Prien am Chiemsee, Germany). The time series of 150 patients (120 females and 30 males) were included in the analysis on the basis of a criterion of less than 20% missing data in a time series. The mean number of missing data in this sample was 2.5 days (=measurement points; standard deviation [SD]: 4.0), which corresponds to a compliance rate of 96.4%. The mean time series length was 69.1 days (SD: 22.6; see Table 1). The inclusion criterion of patients with less than 20% missing data (by this we excluded about 20% of the patients treated during the reference period of about 2 years) is due to the necessity of having time series with high variability (missing data produce straight lines in the process) to get a realistic picture of the dynamics and to get valid inter-item correlations. This procedure is justified because this is not a compliance study, which should identify realistic compliance and missing rates (compare Schiepek, Aichhorn, et al., 2016), but a factor analysis, which requires a sufficient density of time series data.

For 108 out of 150 patients, ICD-based Symptom Rating (ISR)-based assessments at the beginning of the hospital stay (pretreatment) were available (Tritt, 2015; Tritt et al., 2008), which was used for a

TABLE 1 Patient characteristics of the sample included in the factor and item analysis of the Therapy Process Questionnaire

	Total	UHPS	SIP
N	150	105	45
m/f	30/120	27/78	3/42
Age, AM (SD)	41.3 (11.8)	41.71(11.48)	40.22 (12.54)
Time series length (days), AM (SD)	69.1 (22.6)	74.04 (22.45)	57.58 (18.64)
Missing days, AM (SD)	2.5 (4.0)	2.3 (4.1)	2.9 (3.8)
Compliance rate, AM% (SD%)	96.4 (5.7)	96.9 (5.4)	95.1 (6.3)
First-order diagnoses	Total N(%)	UHPS N(%)	SIP N(%)
Bipolar affective disorder (F31.1, F31.3, F31.6, F31.7)	5 (3.3)	5 (4.8)	0
Depressive episode/recurrent depressive disorder (F32.1, F32.2, F32.3, F33.0, F33.1, F33.2, F33.3, F33.4)	40 (26.7)	40 (38.1)	0
Phobic anxiety disorders/other anxiety disorders (F40.1, F41.0, F41.2)	5 (3.3)	5 (4.8)	0
Obsessive-compulsive disorder (F42.0, F42.1, F42.2)	8 (5.3)	8 (7.6)	0
Reaction to severe stress/adjustment disorders (F43.0, F43.1, F43.2)	66 (44.0)	44 (41.9)	22 (48.9)
Dissociative (conversion) disorders (F44.8, F44.81, F44.9)	10 (6.6)	0	10 (22.2)
Somatiform disorders (F54.0, F45.40, F45.41)	3 (2.0)	0	3 (6.7)
Disorders of adult personality and behaviour (F60.30, F60.31, F60.6, F60.8, F61.0)	8 (5.3)	0	8 (17.8)
Other disorders (F10.21, F23.8, F25.1, F48.1, F91.0)	5 (3.3)	3 (2.9)	2 (4.4)

Abbreviations: AM, arithmetic mean; SD, standard deviation; SIP, patients treated at the Department for Psychotraumatology, Clinic St. Irmingard, Prien; UHPS, subsample of patients who were treated at the Department of Inpatient Psychotherapy, University Hospital of Psychiatry, Psychotherapy and Psychosomatics, Salzburg.

construct validation of the subscales of the TPQ (not all patients filled in the ISR pretreatment and posttreatment). The ISR is a first-order criterion outcome measure, assessing symptom severity corresponding to the criteria of the diagnostic F-clusters of the ICD-10. It is not a process monitoring system, but we can expect—and this would be at least a partial construct validation—that some of the factors of the TPQ may be positively or negatively linked to symptom severity; for example, the factors (see below) “well-being and positive emotions” (WPE), “motivation for change” (MOT), or “mindfulness/self-care” (MSC) are supposed to be negatively correlated with symptom severity, whereas “emotional and problem intensity” (EPI) may be positively correlated with symptom severity.

The diagnoses of the patients (following the nomenclature of the ICD-10) are shown in Table 1. The two main diagnostic clusters of the sample are “depressive episode/recurrent depressive disorder” (26.7%) and “reaction to severe stress/adjustment disorders” (44.0%). The mean numbers of co-diagnoses are 0.80 (bipolar affective disorders), 1.13 (depressive episode/recurrent depressive disorder), 0.60 (phobic anxiety disorders/other anxiety disorders), 1.13 (obsessive-compulsive disorder [OCD]), 2.73 (reaction to severe stress/adjustment disorders), 2.17 (dissociative [conversion] disorders), 1.67 (somatoform disorders), 2.88 (disorders of adult personality and behaviour), and 2.00 (other disorders).

Ethical approval for the application of the TPQ by using an internet-based device for patient monitoring and the usage of the retrieved data was given by the ethical committee of the Salzburg County Governance. Daily self-ratings by using the TPQ were based on a written informed consent of all patients. All procedures were in accordance with the Helsinki Declaration as revised in 2013.

2.2 | Questionnaires and monitoring technology

The TPQ was administered in a day-by-day routine during inpatient psychotherapy. For reasons of cross-validation, we used the ISR (Fischer, Tritt, Klapp, & Fliege, 2009, 2011; Tritt, 2015; Tritt et al., 2008), which was applied at the beginning (pre) and at the end (post) of the treatment. The subscales of the ISR are “depression,” “anxiety,” “OCD,” “somatoform disorder,” “eating disorder,” and an additional scale with problems not related to the other subscales. The total score of the ISR averages all subscales by a weight of 1, the additional scale by a weight of 2. Both questionnaires, the TPQ and the ISR, were administered by an internet- or app-based (this is up to free choice) monitoring system, the SNS, which was developed for the assessment of processes and outcomes in naturalistic settings (Schiepek, Aichhorn, et al., 2016; Schiepek, Aichhorn, & Schöller, 2018; Schiepek, Stöger-Schmidinger, et al., 2016).

2.3 | Construction and history of the TPQ

The items of the TPQ were created by a literature research on psychological factors contributing to the engagement of patients in their change process (“process involvement”), a construct close to that of

“self-relatedness,” which is a core concept of the Generic Model of Psychotherapy (Orlinsky, 2009; Orlinsky & Howard, 1986; Orlinsky, Ronnestad, & Willutzki, 2004). The questionnaire was factor analysed on the basis of data from a process–outcome study with inpatients treated in routine practice (Honeremann & Schiepek, 2006; $N = 94$ cases, seven factors with 59.6% explained variance). A second factor analysis was based on another sample of inpatients (Schiepek, Aichhorn, & Strunk, 2012; $N = 149$ cases, five factors with 74.5% explained variance). Beginning from 2007, the TPQ was presented and filled in by an electronic, internet-based device, the SNS (Schiepek et al., 2015; Schiepek, Aichhorn, et al., 2016). After some years of experience in different inpatient settings, the items were revised by the team of the Department of Inpatient Psychotherapy of the Christian Doppler Medical Center Salzburg. On the basis of clinical experience, some items were omitted, others were newly introduced. Items were omitted because of unclear meaning (e.g., “Today, I felt compassion”—to whom, self or others? or “Today I felt confused.”) New items were introduced because mindfulness, self-care, and body-related experiences seemed important to most of the therapists in the team. The hitherto combined 7-point Likert scales and visual analogue scales were replaced by visual analogue scales only for all 47 items.¹ This revised TPQ was used from spring 2016 up to now. The data used for the exploratory factor analysis and confirmatory factor analysis (CFA) were collected during this period.

2.4 | Statistics

The exploratory factor analysis was realized by the methods of principal component analysis (PCA) and principal axis factoring using SPSS 24.0.0.1 (64 Bit version). The adequacy of the data for factor analysis was tested by the Kaiser–Meyer–Olkin (KMO) score and the Bartlett's test of sphericity. The KMO score is a measure of the proportion of variance among the included variables that might be common variance; scores between 0.8 and 1 indicate the appropriateness of the sample for factor analysis. The Bartlett's test checks if the correlation matrix of the included items deviates from the unity matrix, that is, if the correlations are unequal to 0.

The CFA (Brown, 2015) was realized in R (Studio Version 1.1.463; © 2009–2018, RStudio Inc with R-version 3.5.0 [2018-04-23], and R packages “lavaan”[version 0.6–3] and “psych” [version 1.8.10]) using the χ^2 fit criterion, the Tucker–Lewis index (TLI), the comparative fit index (CFI), and the root mean square error of approximation (RMSEA). The TLI (Tucker & Lewis, 1973) analyses the discrepancy between the chi-squared value of the hypothesized model and the null model. This nonnormed fit index resolves some problems of the negative bias of the normed fit index. Its range is between 0 and 1, with a cut-off for values $>.95$ indicating a good model fit (Hu & Bentler,

¹The questionnaire is introduced by the following “welcome statement”: Welcome and thank you for your participation in the therapy monitoring. Please fill in the following questionnaire by rating your experiences today. Some of the items concern your therapeutic progress or the relationship with your therapists and other patients. Even if you did not have direct contact with therapists or fellow patients today, please still try to answer the questions. Please answer according to how your experiences, impressions, and mood made you feel today.

1999). The CFI analyses the model fit by examining the discrepancy between the data and the hypothesized model while adjusting for the issues of sample size inherent in the chi-squared test of model fit and the normed fit index (Bentler, 1990). CFI values range from 0 to 1, with larger values indicating better fit. CFI values $>.95$ can be accepted as an indicator of good fit (Bentler & Bonett, 1980; Hu & Bentler, 1999).

The RMSEA avoids issues of sample size by analysing the discrepancy between the hypothesized model, with optimally chosen parameter estimates, and the population covariance matrix (Hooper, Coughlan, & Mullen, 2008). It ranges from 0 to 1, with smaller values indicating better model fit. A value of $.06$ or less is indicative of acceptable model fit. The standardized root mean square residual (SRMR) indicates the square root of the difference between the residuals of the sample covariance matrix and the hypothesized covariance model (Kline, 2005). Values of the SRMR range from 0 to 1, with models obtaining values less than $.08$ being deemed acceptable (Hu & Bentler, 1999).

The item analysis and psychometric characteristics of the TPQ includes Cronbach's α , a measure of the internal consistency of a scale, the mean intercorrelation between the items of a subscale, and the mean discriminative power, which is the mean correlation of each item of a subscale with the total score of the subscale. The calculation was realized by SPSS 24.0.0.1 (64 Bit version, subprogram "reliability analysis"). An overall calculation of Cronbach's α was realized for the factors (subscales) on the basis of the complete set of linked multiple time series of all items over 10.442 measurement points. A further calculation of Cronbach's α values was realized for each time point. Given the fact that the mean time series length of our sample is 69.1 (the next integer value is 70, see Table 1), we calculated the values of each measurement point from 1 to 70. This procedure ensures that more than 50% of all subjects (85 out of 150) were included in the calculation. In Table 2, the mean Cronbach's α of the measurement points from 1 to 70 and the minimum and maximum values out of 70 calculations are reported. The analysis procedure was realized as proposed by Cronbach (1951) using R version 3.5.3 (R Core Team, 2019) and the ltm package (Rizopoulos, 2006).

3 | RESULTS

3.1 | Factor analysis

The database of the factor analysis was a multiple time series of the 47 items of the TPQ. The time series of all 150 subjects were linked together, resulting in an artificial time series of 10.442 measurement points. The inter-item correlation matrix of all items was calculated on this multiple time series. A KMO score of 0.968 and a highly significant ($\chi^2 = 426,042.12$, degrees of freedom [df] = 903, $p < .001$) Bartlett's test of sphericity confirmed the adequacy of the data for factor analysis, that is, the correlations in the correlation matrix are unequal to 0.

3.1.1 | Exploratory factor analysis

The factor analysis was realized by the methods of PCA and principal axis factoring, with similar results for the number of extracted factors and the assignment of the items to the factors with an assignment criterion of 0.5 for the rotated factor matrix. As the factor loadings obtained by varimax and oblimin direct ($\delta = 0$) rotation yielded similar results, we report on the solutions of the PCA method with varimax rotation. The exploratory factor analysis (PCA) revealed a seven-factor solution, with the criterion of an eigenvalue >1 . The seven-factor solution accounts for 68.68% of the variance of the inter-item correlations. Table 2 shows the eigenvalues of the extracted factors, the variance, which is explained by each factor, and the cumulated variance of all factors. Table 3 presents the items of the TPQ and their association to the factors, the factor loadings of all items on the corresponding factor, the arithmetic mean and the standard deviation of the factor's and item's time series, and the communalities of the items (h^2). All items realize high and distinct factor loadings, indicating exclusive item-factor associations of each item to one specific factor.

Items with lower factor loadings (cut-off criterion: $<.5$) or insufficient distinctness of the loadings between different factors were eliminated from the list of items (Table 4). This concerns the items "Today, I was

TABLE 2 Characteristics of the factors (subscales) of the TPQ

Factor	EV	Var%	VarC%	$C\alpha$	$C\alpha(70)$ (min-max)	IIC	MDP	n
Well-being and positive emotions (WPPE)	6.81	15.84	15.84	0.944	0.922 (0.881-0.945)	0.711	0.816	7
Relationship with fellow patients (RFP)	6.13	14.27	30.11	—	—	0.552	0.552	2
Therapeutic alliance and clinical setting (TAS)	4.81	11.19	41.30	0.944	0.919 (0.855-0.946)	0.740	0.832	6
Emotional and problem intensity (EPI)	4.75	11.05	52.35	0.919	0.904 (0.826-0.935)	0.595	0.717	9
Insight/confidence/therapeutic progress (ICP)	3.47	8.08	60.43	0.943	0.931 (0.854-0.953)	0.625	0.766	10
Motivation for change (MOT)	2.47	5.75	66.17	0.928	0.891 (0.821-0.921)	0.721	0.812	5
Mindfulness/self-care (MSC)	1.08	2.51	68.68	0.924	0.866 (0.789-0.891)	0.753	0.824	4

Abbreviations: $C\alpha$, Cronbach's α based on the complete set of multiple time series of all items over 10.442 measurement points; $C\alpha(70)$, mean of all values of Cronbach's α calculated for the measurement points 1 to 70; EV, eigenvalues of the extracted factors (stop criterion: eigenvalue >1); IIC, mean inter-item correlations of the factors; MDP, mean discriminative power of the items of each factor; (min-max), smallest and largest value of Cronbach's α calculated for the measurement points 1 to 70 (the mean time series length of 70 still includes 85 out of 150 patients); n , number of items of each factor; TPQ, Therapy Process Questionnaire; Var%, explained variance of the factors; VarC%, cumulated explained variance of the factors.

TABLE 3 Arithmetic mean, standard deviation, factor loadings, and communalities (h^2) of the items corresponding to the factors of the TPQ

Item (VAS description)	AM	SD	WPE	RFP	TAS	EPI	ICP	MOT	MSC	h^2
Well-being and positive emotions (WPE)	41.50	24.11								
1 Today I felt joy (not at all – very much)	41.99	28.22	0.748	0.102	0.060	-0.285	0.280	0.139	0.167	0.774
2 Today I experienced moments of happiness and light-heartedness (not at all-very much)	39.48	28.21	0.746	0.083	0.042	-0.311	0.280	0.109	0.147	0.768
3 Today I felt energized (not at all-very much)	40.73	26.65	0.670	0.069	0.064	-0.328	0.337	0.154	0.123	0.770
4 Today my self-esteem was (very low-very high)	38.70	26.74	0.651	0.124	0.041	-0.346	0.348	0.174	0.250	0.703
5 Today I felt comfortable in my body (not at all-very comfortable)	36.53	27.71	0.645	0.113	0.018	-0.352	0.367	0.108	0.175	0.721
6 Today I was satisfied with myself (not at all-very much)	43.12	27.92	0.621	0.080	0.065	-0.373	0.376	0.221	0.262	0.788
7 Today I felt valued (not at all-very much)	49.26	29.20	0.534	0.111	0.053	-0.144	0.305	0.225	0.215	0.501
Relationship with fellow patients (RFP)	65.88	24.25								
8 Today I experienced myself in contact with fellow patients as (not at all tense-very much tense)	67.35	25.92	-0.145	-0.568	-0.034	0.358	-0.038	-0.032	0.002	0.664
9 I feel at ease with other patients (not at all-very much)	35.94	29.30	0.177	0.703	0.214	-0.218	0.163	0.122	0.086	0.479
Therapeutic alliance and clinical setting (TAS)	84.33	17.71								
10 I perceive the work with my therapist(s) as helpful (not at all-very much)	84.28	20.11	0.080	0.033	0.905	-0.047	0.124	0.123	0.053	0.860
11 I feel understood by my therapist(s) (not at all-very much)	84.72	19.49	0.083	0.071	0.889	-0.044	0.080	0.113	0.041	0.824
12 I feel supported by my therapist(s) in my personal development (not at all-very much)	83.90	20.63	0.091	-0.014	0.885	-0.072	0.161	0.061	0.065	0.827
13 I feel safe and supported in the clinic (not at all-very much)	82.48	21.25	0.018	0.121	0.792	-0.100	0.121	0.129	0.039	0.675
14 I can be open and honest with my therapist(s) (not at all-very much)	87.87	17.03	-0.052	0.018	0.784	-0.086	0.169	0.189	0.018	0.627
15 This clinic is the right place for me and my personal development (not at all-very much)	82.65	21.62	0.030	0.020	0.778	-0.053	0.067	0.090	0.050	0.685
Emotional and problem intensity (EPI)	46.60	24.52								
16 Today I felt anxious (not at all-very much)	47.95	32.30	-0.138	-0.056	-0.007	0.814	-0.071	-0.072	-0.098	0.700
17 Today I felt tense and restless (not at all-very much)	58.01	29.94	-0.269	0.009	-0.061	0.792	-0.143	-0.059	0.008	0.667
18 Today my problems/complaints were ... (absent-very intense)	53.93	27.17	-0.171	-0.034	-0.006	0.792	-0.081	-0.027	-0.061	0.723
19 Today I felt impaired by my complaints in my daily routine (not at all-very much)	51.80	28.79	-0.256	-0.001	-0.067	0.775	-0.136	-0.092	0.017	0.690
20 Today I felt sad (not at all-very much)	48.61	33.52	-0.129	-0.036	-0.106	0.743	-0.069	-0.013	-0.061	0.587
21 Today I felt guilty (not at all-very much)	41.18	34.13	-0.090	-0.196	0.037	0.676	0.009	-0.127	-0.180	0.575
22 Today I felt helpless with regard to my problems (not at all-very much)	43.28	30.18	-0.255	-0.001	-0.158	0.662	-0.151	-0.125	-0.092	0.544
23 Today I felt shame (not at all-very much)	39.92	33.50	-0.137	-0.261	-0.027	0.633	-0.015	-0.013	-0.216	0.524
24 Today I felt angry (not at all-very much)	37.30	33.40	-0.017	-0.111	-0.115	0.581	0.004	-0.048	-0.060	0.359
Insight/confidence/therapeutic progress (ICP)	42.43	23.28								
25 Today I became aware of relations that were not clear to me before (not at all-very much)	42.22	28.73	0.153	0.047	0.135	-0.050	0.818	0.150	0.136	0.757
26 I now understand myself and my problems better (not at all-very much)	45.13	27.22	0.208	0.027	0.166	-0.151	0.792	0.168	0.209	0.786
27 Today I came closer to a solution for my problems (not at all-very much)	40.72	27.66	0.298	0.071	0.131	-0.169	0.792	0.213	0.110	0.817
28 Today I had new insights about how to better deal with my life circumstances (not at all-very much)	39.25	27.39	0.305	0.065	0.108	-0.155	0.784	0.171	0.115	0.782

(Continues)

TABLE 3 (Continued)

Item (VAS description)	AM	SD	WPE	RFP	TAS	EPI	ICP	MOT	MSC	h^2
29 Today I gained insight into how my thoughts, feelings, and behaviour influence each other (not at all–very much)	46.30	28.74	0.143	0.010	0.125	−0.133	0.709	0.232	0.174	0.640
30 Today I felt able to deal with situations that I never felt able to deal with before (not at all–very much)	39.70	27.95	0.371	0.067	0.092	−0.191	0.685	0.184	0.140	0.704
31 Today I felt confident to approach burdensome issues in my life (not at all–very much)	46.85	28.76	0.169	0.061	0.068	0.036	0.617	0.270	0.179	0.522
32 Today I worked on things that were new and unusual for me (not at all–very much)	38.84	28.86	0.113	−0.003	0.100	0.133	0.564	0.159	0.016	0.381
33 Today I could take a positive view on myself and my progress (not at all–very much)	46.97	27.95	0.405	0.034	0.151	−0.293	0.530	0.303	0.276	0.718
34 Today I felt confident that I will resolve my issues (not at all–very much)	50.61	27.27	0.336	0.083	0.248	−0.251	0.484	0.415	0.136	0.662
Motivation for change (MOT)	58.88	24.34								
35 Today I felt motivated to work on accomplishing my goals (not at all–very much)	59.97	27.50	0.211	0.042	0.209	−0.116	0.372	0.787	0.108	0.868
36 Today I felt determined to tackle my problems (not at all–very much)	61.02	27.48	0.181	0.047	0.242	−0.121	0.345	0.750	0.153	0.810
37 Today I was committed to accomplish my goals (not at all–very much)	57.41	27.24	0.190	0.037	0.174	−0.083	0.435	0.720	0.132	0.795
38 Today I had my goals clearly in mind (not at all–very much)	54.19	28.42	0.214	0.057	0.201	−0.165	0.420	0.650	0.134	0.727
39 Today my interest in the topics of therapy was (very low–very high)	61.39	27.82	0.080	0.080	0.254	−0.061	0.361	0.515	0.086	0.489
Mindfulness/self-care (MSC)	49.83	23.96								
40 Today I treated myself with care (not at all–very much)	49.34	26.25	0.315	0.016	0.108	−0.241	0.350	0.195	0.702	0.770
41 Today I paid attention to my boundaries/limits (not at all–very much)	49.11	25.94	0.315	0.008	0.094	−0.244	0.321	0.164	0.695	0.816
42 Today I paid attention to my bodily signals (not at all–very much)	49.38	26.87	0.275	0.086	0.101	−0.144	0.336	0.128	0.689	0.711
43 Today I was aware of my own needs (not at all–very much)	51.21	26.81	0.432	0.064	0.081	−0.276	0.362	0.269	0.521	0.743

Note. Factor loadings indicate the item–factor association.

Abbreviations: AM, arithmetic mean of all factors and items, with reference to a scale range from 0 to 100; SD, standard deviation; TPQ, Therapy Process Questionnaire; VAS, visual analogue scales.

TABLE 4 Excluded items

Item	AM	SD	WPE	RFP	TAS	EPI	ICP	MOT	MSC
Today I was able to cope with interpersonal conflicts	50.66	28.12	0.332	0.270	0.073	−0.181	0.229	0.259	0.215
Today I experienced a conflict with the therapist(s) or other members of the team of professionals	7.72	15.82	0.113	−0.213	−0.181	0.200	0.149	−0.067	0.038
Today I avoided situations which were related to my problems	38.02	28.65	−0.053	−0.243	0.036	0.280	0.045	−0.267	0.068
Today I was able to manage my emotions	47.35	25.87	0.470	0.124	0.056	−0.409	0.373	0.232	0.359

Note. Items were excluded from the list of respected items because of low and insufficiently differentiating factor loadings (cut-off: <.5). The table shows the factor loadings and communalities (h^2) of the four excluded items.

Abbreviations: AM, arithmetic mean, with reference to a scale range from 0 to 100; EPI, emotional and problem intensity; ICP, insight/confidence/therapeutic progress; MOT, motivation for change; MSC, mindfulness/self-care; RFP, relationship with fellow patients; SD, standard deviation; TAS, therapeutic alliance and clinical setting; WPE, well-being and positive emotions.

able to cope with interpersonal conflicts,” “Today I experienced a conflict with the therapist(s) or other members of the team of professionals,” and “Today I avoided situations which were related to my problems.” One item was excluded not only because it differentiated

insufficiently, but also because it was reported by the patients as unclear and difficult to understand (“Today I was able to manage my emotions”). After these four items were omitted, the TPQ now comprises 43 items.

TABLE 5 Chi-square and fit indices of the CFA

Measure	Value
χ^2	22,995.72 ($p < .001$)
TLI	0.984
CFI	0.985
RMSEA	0.049
SRMR	0.051

Note. Cut-off scores for good model fit: TLI, $>.95$; CFI, $>.95$; RMSEA, $<.05$; SRMR, $<.06$.

Abbreviations: CFA, confirmatory factor analysis; CFI, comparative fit index; RMSEA, root mean square error of approximation; SRMR, standardized root mean square residual; TLI, Tucker–Lewis Index.

3.1.2 | Confirmatory factor analysis

All fit indices of the CFA (Brown, 2015) confirmed the exploratory factor analysis, as can be seen in Table 5. The χ^2 of the CFA fit is 22,995.72 ($p < .001$). All criteria (relative fit indices) of our CFA (the TLI, CFI, RMSEA, and SRMR) exceed respectively fall below the thresholds of good model fits (Table 5; compare Baumgartner & Homburg, 1996; Hu & Bentler, 1999).

3.1.3 | Interfactor correlations

The intercorrelations between the extracted factors confirm relations that can be expected from psychotherapy research and practical experience (Table 6). The EPI factor is negatively correlated with all other factors, with the most negative correlation ($r = -.57$) realized with the WPE factor. The highest positive correlations are given between WPE and MSC ($r = .72$), between ICP and MOT ($r = .69$), between ICP and WPE ($r = .66$), and between ICP and MSC ($r = .64$). The WPE-RFP correlation is almost twice as high ($r = .40$) as the WPE-TAS correlation ($r = .21$). The negative correlation between EPI and RFP is twice as high as that between EPI and TAS ($r = -.44$ vs. $r = -.18$), but MOT is more highly correlated with TAS ($r = .43$) than with RFP ($r = .27$; for an interpretation of the results, see Section 4).

TABLE 6 Factor intercorrelations

	WPE	RFP	TAS	EPI	ICP	MOT	MSC
Well-being and positive emotions (WPE)		0.40	0.21	-0.57	0.66	0.56	0.72
Relationship with fellow patients (RFP)			0.22	-0.44	0.26	0.27	0.30
Therapeutic alliance and clinical setting (TAS)				-0.18	0.31	0.43	0.26
Emotional and problem intensity (EPI)					-0.28	-0.30	-0.44
Insight/confidence/therapeutic progress (ICP)						0.69	0.64
Motivation for change (MOT)							0.56
Mindfulness/self-care (MSC)							

Note. The correlations are based on the z-transformed time series of the factors. These time series are produced by including all patients of the sample, which resulted in an artificial time series of 10.442 measurement points. All correlations are significant at $p < .001$. The applied family-wise error correction for multiple comparisons did not change the significance levels of the correlations.

3.2 | Item analysis and psychometric data

For each subscale of the TPQ, some important psychometric characteristics were calculated: Cronbach's α , a measure of the internal consistency of a scale, the mean intercorrelations between items, and the mean discriminative power, which is the mean correlation of each item of a subscale with the total score of the subscale. Whereas the internal consistency and the discriminative power should be high, the item intercorrelations should be in a mean range, because the items of a subscale should not identically represent the same feature of an "object." Table 2 presents the results of the item analysis and subscale characteristics.

Concerning the calculation of Cronbach's α , it should be respected that the procedure is based on some restrictive assumptions, that is, unidimensionality, uncorrelated errors, and essentially tau-equivalence (Rammstedt & Beierlein, 2014). On the basis of two items as in the RFP subscale, these assumptions cannot be tested, and, in consequence, no values are reported for RFP in Table 2. Beside the calculation of Cronbach's α based on the complete set of multiple time series of all items over 10.442 measurement points, another calculation was realized for each time point. Because the mean of the time series length of our sample is 70, we calculated the values for 1 to 70 measurement points. In Table 2, the mean Cronbach's α of 70 time series points and the minimum and maximum values out of the 70 calculations are reported.

3.3 | Construct validity

The partial construct validity of the factors (subscales) of the TPQ is demonstrated by the intercorrelations with the subscales of the ISR. Because of the fact that the TPQ is not a measure of symptom severity or of other outcome criteria, the correlations between the subscales of the two questionnaires will only reveal a partial construct validity. However, EPI or, reversely, WPE, may directly correspond to symptom severity, whereas other psychological constructs represented by the subscales of the TPQ may intensify or weaken symptom severity, in other words, may be moderators of

TABLE 7 Correlations between the factors of the TPQ and the subscales of the ISR

Factors	Subscales						
	Dep	Anx	OCD	Som	Eat	Add	Total
WPE	-0.61**	-0.39**	-0.16	-0.17	-0.24*	-0.55**	-0.51**
RFP	-0.06	-0.08	0.03	0.00	0.01	-0.10	-0.02
TAS	-0.28**	-0.06	-0.06	0.09	0.04	-0.25*	-0.10
EPI	0.53**	0.37**	0.32**	0.14	0.20*	0.49**	0.50**
ICP	-0.49**	-0.19	-0.15	-0.04	-0.09	-0.35**	-0.29**
MOT	-0.42**	-0.21*	0.03	-0.07	0.08	-0.30**	-0.21*
MSC	-0.49**	-0.29**	-0.11	-0.2*	-0.10	-0.41**	-0.38**

Note. Intercorrelations between the arithmetic mean of the factor values at the first seven measurement points (beginning of the treatment) and the subscales of the ICD-10 based Symptom Rating (ISR), taken at the first days of hospital stay. Subscales: Dep (depression), Anx (anxiety disorder), OCD (obsessive-compulsive disorder), Som (somatoform disorder), Eat (eating disorder), Add (additional scale), Total (total score).

Abbreviations: EPI, emotional and problem intensity; ICP, insight/confidence/therapeutic progress; MOT, motivation for change; MSC, mindfulness/self-care; RFP, relationship with fellow patients; TAS, therapeutic alliance and clinical setting; TPQ, Therapy Process Questionnaire; WPE, well-being and positive emotions.

psychopathology. The correlations were calculated between the ISR at intake to the hospital and the mean of the first seven values of the time series of the TPQ factors. Given daily measures, seven measurement points correspond to 1 week, which ensures that we take a more stable level of the dynamics than by taking a single day. A single measure always can represent by chance a very high or a very low occurrence, which would distort the result. As shown in Table 7, the subscales depression, anxiety, eating disorder, the additional scale, and the total score of the ISR were significantly negative correlated with WPE and positively correlated with EPI. ICP, MOT, and MSC were negatively correlated with depression, anxiety, eating disorder, the additional scale, and the total score of the ISR. EPI was positively correlated with the OCD and the eating disorder subscale, and the MSC factor correlated negatively with the somatoform disorder subscale of the ISR. It should be noted that symptoms related to OCD, somatoform, and eating disorders were less important for most of the patients of our sample than depression- and anxiety-related symptoms (compare Table 1). Although we outlined no specific hypotheses on the correlations between the RFP and the TAS factors with the subscales of the ISR, these correlations are shown in Table 7. There are no significant correlations of RFP with the ISR subscales but significant negative correlations of the TAS with the depression subscale and the additional subscale of the ISR. More depressed patients may have a poorer relationship with professionals at the beginning of the hospital stay but not with fellow patients.

For each factor of the TPQ, the time series can be calculated by averaging the respective items of the factors or by calculating z-transformed dynamics on the averaged items. Figure 2 shows the evolution of the factors representing the therapeutic process of a




FIGURE 2 Time series of the seven factors of the TPQ, illustrated by a single case (104 measurement points). The time series are calculated by averaging the items associated to each factor and then applying a z-transformation, which shows the time series in units of standard deviations on the y-axis around the mean of zero. This is the way how factor dynamics is presented in the SNS. Usually, the z-transformed factor dynamics shows the shape of a process more pronounced than time series resulting only from averaging items.

patient diagnosed with a complex post-traumatic stress disorder. The synchronized order transition during this psychotherapy after about one third of the process can be seen in different factors, for example, EPI, ICP, MOT, and MSC. Other subscales (e.g., TAS) reveal more stable dynamics after an instable period at the beginning, representing stable boundary conditions of the transient destabilization during the self-organizing process.

4 | DISCUSSION

The factor analysis of the TPQ revealed a seven-factor solution including 43 items distributed on the factors (subscales). Four items were excluded for statistical and clinical reasons. The factors address a broad variety of change mechanisms, which are discussed in the scope of common factors research on psychotherapy ("contextual model," Wampold & Imel, 2015).

One stable and frequently replicated result from this research concerns the importance of the therapeutic alliance as a contributor to change (e.g., Flückiger, del Re, Wampold, & Horvath, 2018; Norcross & Lambert, 2011; Wampold, 2015), which is represented by the factor TAS of the TPQ and, also, though investigated to a minor degree, the RFP. In a clinical context, fellow patients take the role of a social network supporting the patient and his development (e.g., Jörgensen, Römme, & Rundmo, 2009). Although it may seem to represent an artificial setting, the RFP at a hospital ward is a contributor and mediator of change because—among other reasons—intensive social interactions enable social learning (Adler & Stead, 2015). In addition to this, the experience of social relations may be a secondary outcome criterion. Other contributors are positive emotions and well-being (compare the results from positive psychology, for example, Seligman & Csikszentmihalyi, 2000; Sin & Lyubomirsky, 2009), as well as the activation of resources (e.g., Grawe, 2004), represented by the factor WPE. This factor may be seen as a supporting variable for the personal development of a patient and at the same time as an important secondary outcome criterion (Wampold et al., 2017). WPE reinforces change and may be an indicator of successful development. In a theoretical model of therapeutic change (Schiepek et al., 2017), experiences of success and positive emotions (both are covered by the WPE factor) are related to motivation for change, insight, and problem reduction, as well as to the long-term evolution of personality traits (e.g., reduced hopelessness or increased self-efficacy, mindfulness, and competences in emotion regulation; Schöller et al., 2018).

Insight (e.g., Castonguay & Hill, 2007; Grawe, 2004), trust in one's personal development (including experiences of self-efficacy; e.g., Maddux, 2013), and experienced success reinforcing the therapeutic work (e.g., Catty, 2004) and acting as an agent of self-efficacy are well-known mechanisms of change (represented by the factor ICP). Of course, MOT is an intensely investigated factor contributing to therapeutic progress (e.g., Grawe, 2004; Ryan, Lynch, Vansteenkiste, & Deci, 2011). During the last two decades, mindfulness and positive self-regard (represented by the factor MSC) were recognized as important therapeutic mechanisms (Bateman & Fonagy, 2013, 2015) and created

therapy approaches of its own (e.g., Mindfulness-based Stress Reduction; e.g., Grossmann, Niemann, Schmidt, & Walach, 2004). Independent of any specific therapeutic approach, mindfulness and self-regard contribute to personal development and stress-reduction in a more general sense. Finally, symptom severity and problem intensity represent a primary outcome criterion, which, in many cases, for example, in all mood and anxiety disorders, is related to worrying and stressful emotions. Both components, symptom severity and negative emotions, are integrated in the factor EPI. In addition to this, many psychotherapeutic approaches emphasize the role of worrying emotions for personal development, for example, during confrontation with anxiety-provoking or trauma-related personal issues, avoidance reduction, or taking in mind suppressed needs, phantasies, and burdensome reminders. Emotion-focused psychotherapy (Greenberg, 2002) specifically focuses on the experience of emotions as a core mechanism of therapeutic change. The theoretical constructs as represented by the factors of the TPQ reflect the importance of interpersonal experiences for personal development (TAS and RFP), as stressed by the contextual model of psychotherapy and by the frequently replicated findings on the contribution of the therapeutic alliance to the outcome (e.g., Flückiger et al., 2018). The TAS and the RFP factors explain a substantial part of the variance of the TPQ (see Table 2) and are significantly correlated to other factors, as EPI and MOT (see Table 6). Five from seven factors are related to the patient and refer to numerous findings, which underline that the patient plays a substantial (if not the main) role in producing the process and creating the outcome of psychotherapy (Bohart & Tallman, 2010; Duncan, Miller, & Sparks, 2004; Orlinsky et al., 2004; Prochaska, Norcross, & DiClemente, 1994).

Some interesting insight into the mechanisms of change is given by the interfactor correlations. The high positive correlations between ICP and MOT, WPE, and MSC underline the important role of insight and confidence in the progress for all other ingredients and contributors to symptom reduction, positive emotions, and well-being—in other words, to therapeutic success and good outcome. The closer connectedness of WPE to the RFP, compared with the relationship with the professional therapists, underlines that in clinical settings (inpatient treatment), the relationship with the fellow patients is important and even may be more important and supporting than the alliance with the professional therapists (e.g., Jörgensen et al., 2009). A cautious interpretation of the higher MOT-TAS correlation ($r = .43$) compared with the MOT-REP correlation ($r = .27$) may be that motivation for change is more supported by professionals than by fellow patients, whereas fellow patients play a greater role for positive feelings and well-being (TAS-EPI: $r = -.18$ vs. RFP-EPI: $r = -.44$; TAS-WPE: $r = .21$ vs. RFP-WPE: $r = .40$).

The factor structure of the TPQ corresponds to the theoretical constructs (variables) of a mathematical model, which explains some essential features of change dynamics in psychotherapy (Schiepek et al., 2017; Schöller et al., 2018). The variables of this model are "therapeutic success," "motivation for change," "dysphoric emotions," "symptom severity," and "insight." The factors of the TPQ as presented here are not identical to these variables but very similar. It should be noted that the therapeutic alliance factor is also represented in the

model, not as a variable but as one out of four parameters that modulate the shape of the nonlinear functions interconnecting the variables. The discrete iterations produced by a set of five coupled nonlinear difference equations—each equation representing the dependency of a variable on other variables or on itself—may be interpreted as discrete day-by-day measures of a psychotherapeutic process. In consequence, the validation of the model can be realized by time series as produced by the TPQ (Schöller et al., 2019). The combination of data-driven computer simulations and time series data available at “real time” opens the way for further developments in artificial intelligence and process control in psychotherapy.

Equidistant high-frequency measures as realized by the TPQ allow for the application of time series analysis methods, which can be used for the identification of nonstationarities and pattern transitions in human change processes. One of these methods is dynamic complexity, which combines the amplitude, the frequency, and the distribution of the measurement values over the available scaling range (Haken & Schiepek, 2010; Schiepek & Strunk, 2010). It is calculated within a gliding window, which runs over the complete time series. Dynamic complexity may be characteristic for specific dynamics (e.g., the emotional instability of borderline personality disorder) or may be a precursor of phase transitions (critical instability). Another feature of psychopathological dynamics is increased or decreased synchronization of emotions and cognitions, which can be seen in colour-coded inter-item correlation matrices calculated in running windows over multiple time series. Locally increased synchronization also may be a precursor of phase transitions (Haken & Schiepek, 2010; Scheffer et al., 2009). Recurrence plots portray recurrent patterns and transients within a time series in a time \times time diagram (Eckmann, Oliffson Kamphorst, & Ruelle, 1987; Webber & Zbilut, 1994).

The complexity of trajectories of change, which is mirrored by these analysis methods, is reflected by the fact that the trajectories do not move on standard tracks (Schiepek et al., under review). Compared with standard tracks, which are produced by averaging time series of patients with the same diagnosis or similar symptom severity at intake, patterns of change as assessed by the TPQ are much more complex, unpredictable, and chaotic. Deviations from standard tracks do not predict poor outcome (Schiepek et al., under review); on the contrary, they result from order transitions and critical instabilities, which are necessary conditions for success (Haken & Schiepek, 2010).

4.1 | Strengths

The data of our study was taken from naturalistic inpatient settings. Patients with a range of diagnoses, which is representative for inpatient treatment, filled in the TPQ during the complete period of their hospital stay. The compliance rate was high, resulting in a very low percentage of missing data. The factor structure, which could be extracted from the covariance matrix of all items, was clear and easy to interpret. The cumulated explained variance of the factors is

comparatively high, and the item, as well as subscale characteristics (internal consistency, inter-item correlations of the subscales, and discriminative power of the items), may be declared as excellent. The CFA clearly verified the exploratory factor analysis. The items and the factors of the TPQ correspond to broadly accepted and validated constructs on mechanisms of change in psychotherapy.

Other than outcome measures, which usually focus on symptom severity and, by this, are counterproductive for daily assessments because patients continuously would be minded on problems instead of personal development, many items of the TPQ refer to resources, competencies, or experiences, which could drive progress. This makes the continuous high-frequency application of the TPQ an auto-catalysing therapeutic intervention of its own, which should be tested in further studies.

4.2 | Limitations

The factor analysis was based on the time series data of 150 patients. Of course, the sample size could have been larger, and, in consequence, a replication study based on a bigger sample of not only inpatients but also outpatients should be realized. This is intended and realistic because of the fact that the TPQ is applied in different hospitals, treatment centres, and therapeutic practices. A disadvantage may be seen in the unequal assignment of the items to the factors, with a range from two items in the RFP factor and 10 items in the ICP factor. However, this is an empirical and statistical result, which has to be recognized and cannot be manipulated. In the strictest sense, the results potentially do not apply to the English translation we present in the manuscript but to the German version of the TPQ. Consequently, future psychometric investigations of the English version of the TPQ are of importance and should be realized by an independent replication and validation study.

A 43-item self-report for daily assessment from one perspective may be too labour-intensive; from another perspective, it would be not intensive enough to match the expectations of “high-frequency” monitoring. However, years of experience in different clinical settings proved that patients not only accept this investment of time but also appreciate it as a quiet moment of self-reflection, mentalization, and focusing their own personal development. On the other hand, higher frequencies of self-reports (e.g., by event sampling) would interrupt the day, focus the attention too much on the task of self-rating, and finally produce more missing data in the case of losing one day of feedback. Given the claim of a full assessment of a psychotherapeutic process, daily measures have revealed to be a good compromise.

4.3 | Perspectives

The TPQ is not restricted to specific disorders or treatment approaches. None of the items picks up specific symptoms of specific diagnoses. On the contrary, the questionnaire addresses different psychological constructs and general mechanisms of change. In consequence, it is not only a transdiagnostic measure but also independent

from specific therapeutic concepts, techniques, or psychotherapeutic confessions. As experience shows, therapists and treatment centres working with completely different methods can use the TPQ-based process monitoring.

The use of the TPQ in outpatient settings is possible if the items representing the RFP (evidently, fellow patients do not exist in outpatient psychotherapy) and representing the therapeutic alliance will be omitted. Outpatient psychotherapy usually has a rhythm of weekly or bi-weekly sessions, and it is useless to ask for the quality of the professional relationship each day. All other items can be applied to the monitoring of outpatient psychotherapy.

The TPQ standardizes the sampling rate because it imposes self-assessments at a daily rhythm. This is the reason why the TPQ can be used in very different inpatient, day-treatment, or outpatient settings for routine monitoring and, by this, allows for a direct comparison of very different treatment cases and processes. Its usability not only in studies but in everyday routine practice creates big data sets, which can be analysed for features of the dynamics and their relation to the outcome. Process features such as sudden gains or losses, other kinds of dynamic or order transitions, crisis-repair sequences, critical instabilities, changing synchronization patterns, or early drop-outs can be assessed and even defined in a general way. The definition, assessment, and analysis of such phenomena requires a standardized high-frequency sampling rate.

The option of a differentiated access to dynamic patterns not only allows for new target concepts in diagnostics, such as the diagnostics of stability or instability of the process, of dynamic connectivity and transient synchronization patterns, or of precursors of critical transitions, but also for more general definitions of outcome. Traditionally, we focus on pre-post measures of primary or secondary outcome criteria, which can now be complemented by criteria of changed dynamic patterns, for example, reduced pathological oversynchronization of cognitions and emotions (e.g., Schiepek, Stöger-Schmidinger, et al., 2016), increased flexibility and adaptability of psychological processes (Kashdan & Rottenberg, 2010), or modified emotional dynamics, for example, from emotional instability to stability (e.g., in borderline personality disorder), or the other way round, from emotional rigidity to flexibility (e.g., in major depressive disorder, alexithymia, or addictive disorders).

As we noted, the TPQ was developed for applications in routine real-time monitoring by electronic devices, using internet and app-based systems (e.g., the SNS). By this, it can be seen as an important step towards the digitalization of psychotherapy. In psychotherapy 4.0, computer-assisted methods open new ways for treatment control, just in time applications of interventions, combining web-based and face-to-face treatments, or integrated care linking treatments over different settings (e.g., inpatient-outpatient). Certainly, psychotherapy 4.0 will not be the solution for everything, but combined with new developments in artificial intelligence, it will contribute to the development of useful concepts and help to bridge challenging gaps, for example, between personalized treatment and big data, idiographic and nomothetic approaches, or process control and the personal responsibility of patients.

ACKNOWLEDGEMENTS

We are grateful to David Pincus, PhD, (Chapman University) for a detailed check of the wording of the items that were translated from German to English language.

AUTHORS' CONTRIBUTIONS

G.S. developed the original version of the TPQ, realized the revision together with H.K. and B.S.-S., wrote the paper, and supervised the study and the statistical procedures. H.S. and L.K. realized the statistical analysis. K.V. supervised and checked the statistical procedures. A. L.-A. and G.S. translated the items into English language. L.K., P.H., W. A., B.S.-S., and H.K., together with their teams, were responsible for the data collection in different treatment centres.

ORCID

Kathrin Viol  <https://orcid.org/0000-0003-3482-801X>

Anna Lichtwarck-Aschoff  <https://orcid.org/0000-0002-4365-1538>

REFERENCES

- Adler, N. E., & Stead, W. W. (2015). Patients in context—EHR capture of social and behavioral determinants of health. *New England Journal of Medicine*, *372*, 698–701. <https://doi.org/10.1056/NEJMp1413945>
- Alexander, L. B., & Luborsky, L. (1986). The Penn Helping Alliance Scales. In L. S. Greenberg, & W. M. Pinsof (Eds.), *The psychotherapeutic process: A research handbook* (pp. 325–366). New York, NY: Guilford Press.
- Bateman, A., & Fonagy, P. (2013). Mentalization-based treatment. *Psychoanalytic Inquiry*, *33*, 595–613. <https://doi.org/10.1080/07351690.2013.835170>
- Bateman, A., & Fonagy, P. (2015). Borderline personality disorder and mood disorders: Mentalizing as a framework for integrated treatment. *Journal of Clinical Psychology*, *71*, 792–804. <https://doi.org/10.1002/jclp.22206>
- Baumgartner, H., & Homburg, C. (1996). Applications of structural equation modelling in marketing and consumer research: A review. *International Journal of Research in Marketing*, *13*, 139–161. [https://doi.org/10.1016/0167-8116\(95\)00038-0](https://doi.org/10.1016/0167-8116(95)00038-0)
- Bech, P., Gudex, C., & Johansen, K. S. (1996). The WHO Well-being Index: Validation in diabetes. *Psychotherapy and Psychosomatics*, *65*, 183–190. <https://doi.org/10.1159/000289073>
- Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin*, *107*(2), 238–246. <https://doi.org/10.1037/0033-2909.107.2.238>
- Bentler, P. M., & Bonett, D. G. (1980). Significance tests and goodness of fit in the analysis of covariance structures. *Psychological Bulletin*, *88*, 588–606. <https://doi.org/10.1037/0033-2909.88.3.588>
- Bohart, A. C., & Tallman, K. (2010). Clients: The neglected common factor in psychotherapy. In B. Duncan, S. Miller, B. Wampold, & M. Hubble (Eds.), *2010The heart and soul of change* (2nd. ed.) (pp. 83–111). Washington DC: American Psychological Association.
- Brattland, H., Koksvik, J. M., Burkeland, O., Gråwe, R. W., Klöckner, C., Linaker, O. M., ... Iversen, V. C. (2018). The effects of routine outcome monitoring (ROM) on therapy outcomes in the course of an implementation process: A randomized clinical trial. *Journal of Counseling Psychology*, *65*(5), 641–652. <https://doi.org/10.1037/cou0000286>
- Brockmann, J., Kirsch, H., Hatcher, R., Andreas, S., Benz, S., & Sammet, I. (2011). Dimensionen der therapeutischen Beziehung aus Patienten-

- Perspektive – Entwicklung der "Skala Therapeutische Allianz Revised STA-R". [Dimensions of the therapeutic alliance from the patient's perspective—Development of the "Scale of the Therapeutic Alliance-Revised (STA-R)"]. *Psychotherapie, Psychosomatik, Medizinische Psychologie*, 61, 208–215. <https://doi.org/10.1055/s-0030-1263142>
- Brown, T. (2015). *Confirmatory factor analysis for applied research*. New York: Guilford Press.
- Castonguay, L. G., & Hill, C. (Eds.) (2007). *Insight in psychotherapy*. Washington, DC: American Psychological Association. <https://doi.org/10.1037/11532-000>
- Catty, J. (2004). The vehicle of success': Theoretical and empirical perspectives on the therapeutic alliance in psychotherapy and psychiatry. *Psychology and Psychotherapy*, 77(2), 255–272. <https://doi.org/10.1348/147608304323112528>
- Cronbach, L. J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika*, 16, 297–334. <https://doi.org/10.1007/BF02310555>
- de Jong, K., Barkham, M., Wolpert, M., & the INSPIRE Research Network (in prep). *The impact of progress feedback on outcomes of psychological interventions: A review of the evidence and proposal for a way forward for research*. NL: Manuscript, Institute of Psychology, Leiden University.
- de Jong, K., Segaar, J., Ingenhoven, T., van Busschbach, J., & Timman, R. (2018). Adverse effects of outcome monitoring feedback in patients with personality disorders: A randomized controlled trial in day treatment and inpatient settings. *Journal of Personality Disorders*, 32, 393–413. https://doi.org/10.1521/pedi_2017_31_297
- de Jong, K., van Sluis, P., Nugter, M. A., Heiser, W. J., & Spinhoven, P. (2012). Understanding the differential impact of outcome monitoring: Therapist variables that moderate feedback effects in a randomized clinical trial. *Psychotherapy Research*, 22, 464–474. <https://doi.org/10.2080/10503307.2012.673023>
- Delgadillo, J., Overend, K., Lucock, M., Groom, M., Kirby, N., McMillan, D., ... de Jong, K. (2017). Improving the efficiency of psychological treatment using outcome feedback technology. *Behavior Research and Therapy*, 99, 89–97. <https://doi.org/10.1016/j.brat.2017.09.011>
- Duncan, B., Miller, S., & Sparks, J. A. (Eds.) (2004). *The heroic client: A revolutionary way to improve effectiveness through client-directed, outcome-informed therapy*. San Francisco: Jossey-Bass.
- Duncan, B., Miller, S., Wampold, B., & Hubble, M. (Eds.) (2010). *The heart and soul of change* (2nd ed.). Washington: APA.
- Eckmann, J. P., Oliffson Kamphorst, S., & Ruelle, D. (1987). Recurrence plots of dynamical systems. *Europhysics Letters*, 4, 973–977. <https://doi.org/10.1209/0295-5075/4/9/004>
- Elliott, R., & Wexler, M. M. (1994). Measuring the impact of sessions in process-experiential therapy of depression: The Session Impacts Scale. *Journal of Counseling Psychology*, 41, 166–174. <https://doi.org/10.1037/0022-0167.41.2.166>
- Finch, A. E., Lambert, M. J., & Schaalje, B. G. (2001). Psychotherapy quality control: The statistical generation of expected recovery curves for integration into an early warning system. *Clinical Psychology & Psychotherapy*, 8, 231–242. <https://doi.org/10.1002/cpp.286>
- Fischer, H. F., Tritt, K., Klapp, B. F., & Fliege, H. (2009). Faktorstruktur und psychometrische Eigenschaften des ICD-10-Symptom-Rating (ISR) an Stichproben psychosomatischer Patienten [Factor structure and psychometric data of the ICD-10-based Symptom Rating Scale: Results from samples of psychosomatic patients]. *Psychotherapie, Psychosomatik & Medizinische Psychologie (Psychotherapy, Psychosomatics, & Medical Psychology)*, 60(8), 307–315. <https://doi.org/10.1055/s-0029-1214419>
- Fischer, H. F., Tritt, K., Klapp, B. F., & Fliege, H. (2011). How to compare scores from different depression scales: Equating the Patient Health Questionnaire (PHQ) and the ICD-10-Symptom Rating (ISR) using item response theory. *International Journal of Methods in Psychiatric Research*, 20(4), 203–214. <https://doi.org/10.1002/mpr.350>
- Flückiger, C., del Re, A. C., Wampold, B. E., & Horvath, A. O. (2018). The alliance in adult psychotherapy: A meta-analytic synthesis. *Psychotherapy*, 55(4), 316–340. <https://doi.org/10.1037/pst0000172>
- Flückiger, C., Regli, D., Zwahlen, D., Hostettler, S., & Caspar, F. (2010). Der Berner Patienten- und Therapeutenstundenbogen 2000: Ein Instrument zur Erfassung von Therapieprozessen [The Berne Post Session Report for patients and for therapists 2000: An instrument to measure therapy processes]. *Zeitschrift für Klinische Psychologie und Psychotherapie: Forschung und Praxis*, 39, 71–79.
- Gaston, L. (1991). Reliability and criterion-related validity of the California Psychotherapy Alliance Scales—Patient version. *Psychological Assessment: A Journal of Consulting and Clinical Psychology*, 31, 68–74. <https://doi.org/10.1037/1040-3590.3.1.68>
- Gelo, O. C. G., & Salvatore, S. (2016). A dynamic systems approach to psychotherapy: A meta-theoretical framework for explaining psychotherapy change processes. *Journal of Counseling Psychology*, 63, 379–395. <https://doi.org/10.1037/cou0000150>
- Gibbons, M. B. C., Kurtz, J. E., Thompson, D. L., Mack, R. A., Lee, J. K., Rothbard, A., ... Crits-Christoph, P. (2015). The effectiveness of clinician feedback in the treatment of depression in the community mental health system. *Journal of Consulting and Clinical Psychology*, 83, 748–759. <https://doi.org/10.1037/a0039302>
- Grawe, K. (2004). *Psychological therapy*. Cambridge, MA: Hogrefe.
- Greenberg, L. S. (2002). *Emotion-focused therapy: Coaching clients to work through their feelings*. Washington, DC: American Psychological Association.
- Grossmann, P., Niemann, L., Schmidt, S., & Walach, H. (2004). Mindfulness-based stress reduction and health benefits: A meta-analysis. *Journal of Psychosomatic Research*, 57(1), 35–43. [https://doi.org/10.1016/S0022-3999\(03\)00573-7](https://doi.org/10.1016/S0022-3999(03)00573-7)
- Gumz, A., Brähler, E., Geyer, M., & Erices, R. (2012). Crisis-repair sequences—Considerations on the assessment and classification of breaches in the therapeutic relationship. *BMC Medical Research Methodology*, 12, 10. <https://doi.org/10.1186/1471-2288-12-10>
- Haken, H., & Schiepek, G. (2010). *Synergetik in der Psychologie. Selbstorganisation verstehen und gestalten [Synergetics in psychology. Understanding and supporting self-organization]* (2nd ed., 1st ed. 2006). Göttingen: Hogrefe.
- Hatcher, R.L. & Shannon, P.A. (2005). The patient's collaborative alliance in therapy. Unpublished manuscript.
- Hayes, A. M., Laurenceau, J. P., Feldman, G. C., Strauss, J. L., & Cardaciotto, L. A. (2007). Change is not always linear: The study of nonlinear and discontinuous patterns of change in psychotherapy. *Clinical Psychology Review*, 27, 715–723. <https://doi.org/10.1016/j.cpr.2007.01.008>
- Honermann, H., & Schiepek, G. (2006). The Therapy Process Questionnaire (TPQ). In H. Haken, & G. Schiepek (Eds.), *2006Synergetik in der Psychologie. Selbstorganisation verstehen und gestalten [Synergetics in psychology. Understanding and supporting self-organization]* (2nd. Ed. 2010 ed.) (pp. 363–376). Göttingen: Hogrefe.
- Hooper, D., Coughlan, J., & Mullen, M. R. (2008). Structural equation modelling: Guidelines for determining model fit. *Journal of Business Research Methods*, 6, 53–60.
- Horvath, A. O., & Greenberg, L. S. (1986). The development of the Working Alliance Inventory. In L. S. Greenberg, & W. M. Pinsof (Eds.), *The psychotherapeutic process: A research handbook* (pp. 529–556). New York NY: Guilford Press.
- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives.

- Structural Equation Modeling*, 6(1), 1–55. <https://doi.org/10.1080/10705519909540118>
- Jørgensen, K. N., Rømma, V., & Rundmo, T. (2009). Associations between ward atmosphere, patient satisfaction, and outcome. *Journal of Psychiatric and Mental Health Nursing*, 16(2), 113–120. <https://doi.org/10.1111/j.1365-2850.2008.01333.x>
- Kashdan, T. B., & Rottenberg, J. (2010). Psychological flexibility as a fundamental aspect of health. *Clinical Psychology Review*, 30, 865–878. <https://doi.org/10.1016/j.cpr.2010.03.001>
- Kendrick, T., El-Gohary, M., Stuart, B., Gilbody, S., Churchill, R., Aiken, L., ... Moore, M. (2016). Routine use of patient reported outcome measures (PROMs) for improving treatment of common mental health disorders in adults. *Cochrane Database of Systematic Reviews*, 2016, Issue(7). Art. No), CD011119. <https://doi.org/10.1002/14651858.CD011119.pub2>
- Kline, R. B. (2005). *Principles and practice of structural equation modeling* (3rd ed.). New York: Guilford Press.
- Knaup, C., Kösters, M., Schöfer, D., Becker, T., & Puschner, B. (2009). Effect of feedback of treatment outcome in specialist mental healthcare: Meta-analysis. *The British Journal of Psychiatry*, 195, 15–22. <https://doi.org/10.1192/bjp.bp.108.053967>
- Kratzer, L., Heinz, P., Pfitzer, F., Schennach, R., Aichhorn, W., Aas, B., & Schiepek, G. (2018). Real-time-monitoring in der Behandlung komplexer posttraumatischer Belastungsstörung: Ein Fallbericht [Real-time monitoring in the treatment of complex posttraumatic stress disorder: A case report]. *Verhaltenstherapie*, 28, 93–99. <https://doi.org/10.1159/000481802>
- Lambert, M. J. (2007). Presidential address: What we have learned from a decade of research aimed at improving psychotherapy outcome in routine care. *Psychotherapy Research*, 17, 1–14. <https://doi.org/10.1080/10503300601032506>
- Lambert, M. J. (2010). Yes, it is time for clinicians to routinely monitor treatment outcome. In B. Duncan, S. Miller, B. Wampold, & M. Hubble (Eds.), *The heart and soul of change* (2nd. ed.) (pp. 237–266). Washington DC: American Psychological Association.
- Lambert, M. J. (2017). Maximizing psychotherapy outcome beyond evidence-based medicine. *Psychotherapy and Psychosomatics*, 86, 80–89. <https://doi.org/10.1159/000455170>
- Lambert, M. J., Morton, J. J., Hatfield, D. R., Harmon, C., Hamilton, S., & Shimokawa, K. (2004). *Administration and scoring manual for the OQ-45.2 (Outcome Questionnaire)* (3rd. ed.). Wilmington, DE: American Professional Credentialing Services.
- Lambert, M. J., Whipple, J. L., & Kleinstäuber, M. (2018). Collecting and delivering progress feedback: A meta-analysis of routine outcome monitoring. *Psychotherapy*, 55(4), 520–537. <https://doi.org/10.1037/pst0000167>
- Li, M. J., Black, D. S., & Garland, E. L. (2015). The Applied Mindfulness Process Scale (AMPS): A process measure for evaluating mindfulness-based interventions. *Personality and Individual Differences*, 93, 6–15. <https://doi.org/10.1006/j.paid.2015.10.027>
- Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the Depression Anxiety Stress Scales*. Sydney: Psychology Foundation.
- Lutz, W., & Böhnke, J. R. (2008). Der "Fragebogen zur Evaluation von Psychotherapieverläufen"(FEP-2): Validierungen und Manual [Questionnaire for the evaluation of psychotherapeutic processes: Validation and manual]. *Trierer Psychologische Berichte [Trier Reports in Psychology]*, 35(3).
- Lutz, W., Ehrlich, T., Rubel, J., Hallwachs, N., Röttger, M. A., Joras, C., ... Tschitsaz-Stucki, A. (2013). The ups and downs of psychotherapy: Sudden gains and sudden losses identified with session reports. *Psychotherapy Research*, 23, 14–24. <https://doi.org/10.1080/10503307.2012.693837>
- Maddux, J. E. (2013). *Self-efficacy, adaptation, and adjustment: Theory, research, and application*. New York: Springer Science & Business Media.
- Mander, J., Schlarb, A., Teufel, M., Keller, F., Hautzinger, M., Zipfel, S., ... Sammet, I. (2014). The Individual Therapy Process Questionnaire: Development and validation of a revised measure to evaluate general change mechanisms in psychotherapy. *Clinical Psychology & Psychotherapy*, 22(4), 328–345. <https://doi.org/10.1002/cpp.1892>
- Mander, J., Wittorf, A., Schlarb, A., Hautzinger, M., Zipfel, S., & Sammet, I. (2013). Change mechanisms in psychotherapy: Multiperspective assessment and relation to outcome. *Psychotherapy Research*, 23, 105–116. <https://doi.org/10.1080/10503307.2012.744111>
- Molenaar, P. C. M. (2004). A manifesto on psychology as idiographic science: Bringing the person back into scientific psychology, this time forever. *Measurement*, 2(4), 201–218.
- Molenaar, P. C. M. (2007). Psychological methodology will change profoundly due to the necessity to focus on intra-individual variation. *Integrative Psychological and Behavioral Science*, 41(1), 35–40. <https://doi.org/10.1007/s12124-007-9011-1>
- Molenaar, P. C. M., Sinclair, K. O., Rovine, M. J., Ram, N., & Corneal, S. E. (2009). Analyzing developmental processes on an individual level using nonstationary time series modeling. *Developmental Psychology*, 45(1), 260–271. <https://doi.org/10.1037/a0014170>
- Nilges, P., & Essau, C. (2015). Die Depressions-Angst-Stress-Skalen. Der DASS – ein Screeningverfahren nicht nur für Schmerzpatienten. *Der Schmerz*, 29, 649–657. <https://doi.org/10.1007/s00482-015-0019-z>
- Nischk, D., Grothe, A., & Schiepek, G. (2000). Beratungsprozesse aus Sicht der Klienten. Das mehrdimensionale Konstrukt der Prozessinvolviertheit als Grundlage für die Entwicklung eines Klientenstundenbogens (Consulting processes from the perspective of clients. The multidimensional construct of change involvement as a framework for the development of a session rating scale). *Zeitschrift für Klinische Psychologie, Psychiatrie und Psychotherapie (Journal of Clinical Psychology, Psychiatry, and Psychotherapy)*, 48(2), 118–134.
- Norcross, J. C., & Lambert, M. J. (Eds.) (2011). Psychotherapy relationships that work II. *Psychotherapy*, 48(1), 4–8. <https://doi.org/10.1037/a0022180>
- Orlinsky, D. E. (2009). The "Generic Model of Psychotherapy" after 25 years: Evolution and of a research-based metatheory. *Journal of Psychotherapy Integration*, 19, 319–339. <https://doi.org/10.1037/a0017973>
- Orlinsky, D. E., & Howard, K. I. (1967). The good therapy hour: Experiential correlates of patients' and therapists' evaluations of therapy sessions. *Archives of General Psychiatry*, 16(5), 621–632. <https://doi.org/10.1001/archpsyc.1967.01730230105013>
- Orlinsky, D. E., & Howard, K. J. (1986). Process and outcome in psychotherapy. In S. L. Garfield, & A. E. Bergin (Eds.), *Handbook of psychotherapy and behavior change* (3rd ed.) (pp. 311–381). New York: Wiley.
- Orlinsky, D. E., Ronnestad, M. H., & Willutzki, U. (2004). Fifty years of psychotherapy process-outcome research: Continuity and change. In M. J. Lambert (Ed.), *Bergin and Garfield's handbook of psychotherapy and behavior change* (pp. 307–390). New York: Wiley.
- Patzig, J., & Schiepek, G. (2015). Emotionsregulation und emotionsfokussiertes Prozessmonitoring in der Suchttherapie [Emotion regulation and emotion-focused process monitoring in psychotherapy of chronic alcoholic patients]. In I. Sammet, G. Dammann, & G. Schiepek (Eds.), *HrsgDer psychotherapeutische Prozess [The psychotherapeutic process]* (pp. 124–139). Stuttgart: Kohlhammer.

- Piccirillo, M. L., Beck, E. D., & Rodebaugh, T. L. (2019). A clinician's primer for idiographic research: Considerations and recommendations. *Behavior Therapy*. <https://doi.org/10.1016/j.beth.2019.02.002>
- Prochaska, J. O., Norcross, J. C., & DiClemente, C. C. (1994). *Changing for good*. New York: Morrow.
- R Core Team (2019). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rammstedt, B., & Beierlein, C. (2014). Can't we make it any shorter? The limits of personality assessment and ways to overcome them. *Journal of Individual Differences*, 35(4), 212–220. <https://doi.org/10.1027/1614-0001/a000141>
- Rizopoulos, D. (2006). ltm: An R package for latent variable modelling and item response theory analyses. *Journal of Statistical Software*, 17(5), 1–25. <http://www.jstatsoft.org/v17/i05/>
- Ryan, R. M., Lynch, M. F., Vansteenkiste, M., & Deci, E. L. (2011). Motivation and autonomy in counseling, psychotherapy, and behavior change: A look at theory and practice. *The Counseling Psychologist*, 39(2), 193–260.
- Scheffer, M., Bascompte, J., Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V., ... Sugihara, G. (2009). Early-warning signals for critical transitions. *Nature*, 461, 53–59. <https://doi.org/10.1038/nature08227>
- Schiepek, G., Aichhorn, W., Gruber, M., Strunk, G., Bachler, E., & Aas, B. (2016). Real-time monitoring of psychotherapeutic processes: Concept and compliance. *Frontiers in Psychology for Clinical Settings*, 7, 604. <https://doi.org/10.3389/fpsyg.2016.00604>
- Schiepek, G., Aichhorn, W., & Schöller, H. (2018). Monitoring change dynamics—A nonlinear approach to psychotherapy feedback. *Chaos & Complexity Letters*, 11(3), 355–375.
- Schiepek, G., Aichhorn, W., Schöller, H., & Kronberger, H. (2018). Prozessfeedback in der Psychotherapie. Methodik, Visualisierung und Fallbeispiel [Process feedback on psychotherapy. Methodology, visualization, and case study]. *Psychotherapeut*, 63(4), 306–314. <https://doi.org/10.1007/s00278-018-0272-6>
- Schiepek, G., Aichhorn, W., & Strunk, G. (2012). Der Therapie-Prozessbogen (TPB). Faktorenstruktur und psychometrische Daten [The Therapy Process Questionnaire [TPQ]. Factor structure and psychometric data]. *Zeitschrift für Psychosomatische Medizin und Psychotherapie [Journal of Psychosomatic Medicine and Psychotherapy]*, 58, 257–266. <https://doi.org/10.13109/zptm.2012.58.3.257>
- Schiepek, G., Eckert, H., Aas, B., Wallot, S., & Wallot, A. (2015). *Integrative psychotherapy. A feedback-driven dynamic systems approach*. Boston, MA: Hogrefe International Publishing.
- Schiepek, G., Gelo, O., Viol, K., Kratzer, L., Orsucci, F., de Felice, G., ... Schöller, H. (under review). "Not On Track"—Are psychotherapies moving on standard tracks? *Psychology and Psychotherapy: Theory, Research and Practice*.
- Schiepek, G., Heinzl, S., Karch, S., Plöderl, M., & Strunk, G. (2016). Synergetics in psychology: Patterns and pattern transitions in human change processes. In A. Pelster, & G. Wunner (Eds.), *Selforganization in complex systems: The past, present, and future of Synergetics*. Springer series understanding complex systems. (pp. 181–208). Berlin Heidelberg: Springer.
- Schiepek, G., & Pincus, D. (under review). Complexity science: A framework for psychotherapy integration. *Psychotherapy*.
- Schiepek, G., Stöger-Schmidinger, B., Aichhorn, W., Schöller, H., & Aas, B. (2016). Systemic case formulation, individualized process monitoring, and state dynamics in a case of dissociative identity disorder. *Frontiers in Psychology for Clinical Settings*, 7, 1545. <https://doi.org/10.3389/fpsyg.2016.01545>
- Schiepek, G., & Strunk, G. (2010). The identification of critical fluctuations and phase transitions in short term and coarse-grained time series—A method for the real-time monitoring of human change processes. *Biological Cybernetics*, 102(3), 197–207. <https://doi.org/10.1007/s00422-009-0362-1>
- Schiepek, G., Tominschek, I., & Heinzl, S. (2014). Self-organization in psychotherapy—Testing the synergetic model of change processes. *Frontiers in Psychology for Clinical Settings*, 5/Article 1089, 1–11. <https://doi.org/10.3389/fpsyg.2014.01089>
- Schiepek, G., Viol, K., Aichhorn, W., Hütt, M. T., Sungler, K., Pincus, D., & Schöller, H. (2017). Psychotherapy is chaotic—(not only) in a computational world. *Frontiers in Psychology for Clinical Settings*, 8, 379. <https://doi.org/10.3389/fpsyg.2017.00379>
- Schöller, H., Viol, K., Aichhorn, W., Hütt, M. T., Aas, B., & Schiepek, G. (2018). Personality development in psychotherapy: A synergetic model of state-trait dynamics. *Cognitive Neurodynamics*, 12, 441–459. <https://doi.org/10.1007/s11571-018-9488-y>
- Schöller, H., Viol, K., Goditsch, H., Aichhorn, W., Hütt, M. T., & Schiepek, G. (2019). A nonlinear dynamic systems model of psychotherapy: First steps toward validation and the role of external input. *Nonlinear Dynamics, Psychology, and Life Sciences*, 23(1), 79–112.
- Seligman, M. E. P., & Csikszentmihalyi, M. (2000). Positive psychology: An introduction. *American Psychologist*, 55, 5–14. <https://doi.org/10.1037/0003-066X.55.1.5>
- Shannon, C. E. (1949). Communication in the presence of noise. *Proceedings of the Institute of Radio Engineers*, 37(1). No. 1. (reprint in *Proc. IEEE*, 86(2), 1998)
- Shimokawa, K., Lambert, M. J., & Smart, D. W. (2010). Enhancing treatment outcome of patients at risk of treatment failure: Meta-analytic and mega-analytic review of a psychotherapy quality assurance system. *Journal of Consulting and Clinical Psychology*, 78, 298–311. <https://doi.org/10.1037/a0019247>
- Sin, N. L., & Lyubomirsky, S. (2009). Enhancing well-being and alleviating depressive symptoms with positive psychology interventions: A practice-friendly meta-analysis. *Journal of Clinical Psychology*, 65(5), 467–487. <https://doi.org/10.1002/jclp.20593>
- Smith, S., Hilsenroth, M. J., Baity, M. R., & Knowles, E. S. (2003). Assessment of patient and therapist perspectives of process: A revision of the Vanderbilt Psychotherapy Process Scale. *American Journal of Psychotherapy*, 57(2), 195–205. <https://doi.org/10.1176/appi.psychotherapy.2003.57.2.195>
- Stiles, W. B. (1980). Measurement of the impact of psychotherapy sessions. *Journal of Consulting and Clinical Psychology*, 48(2), 176–185. <https://doi.org/10.1037/0022-006X.48.2.176>
- Stiles, W. B., Glick, M. J., Osatuke, K., Hardy, G. E., Shapiro, D. A., Agnew-Davies, R., ... Barkham, A. (2004). Patterns of alliance development and the rupture-repair hypothesis: Are productive relationships U-shaped or V-shaped? *Journal of Counseling Psychology*, 51, 81–91. <https://doi.org/10.1037/0022-0167.51.1.81>
- Stiles, W. B., Leach, C., Barkham, M., Lucock, M., Iveson, S., Shapiro, D., ... Hardy, G. (2003). Early sudden gains in psychotherapy under routine clinic conditions: Practice-based evidence. *Journal of Consulting and Clinical Psychology*, 71, 14–21. <https://doi.org/10.1037/0022-006X.71.1.14>
- Stiles, W. B., Reynolds, S., Hardy, G. E., Rees, A., Barkham, M., & Shapiro, D. A. (1994). Evaluation and description of psychotherapy sessions by clients using the Session Evaluation Questionnaire and the Session Impact Scale. *Journal of Counseling Psychology*, 41(2), 175–185. <https://doi.org/10.1037/0022-0167.41.2.175>
- Strunk, G., & Schiepek, G. (2006). *Systemische Psychologie. Einführung in die komplexen Grundlagen menschlichen Verhaltens. [Systemic psychology. An*

- introduction to complexity in human behavior and cognition]. München: Elsevier.
- Tritt, K. (2015). *ICD-10-Symptom-Rating (ISR) - das Handbuch zum Fragebogen (Handbook of the ICD-10-based Symptom-Rating [ISR])*. Berlin: Neobooks.
- Tritt, K., von Heymann, F., Zaudig, M., Zacharias, I., Söllner, W., & Löw, T. (2008). Entwicklung des Fragebogens "ICD-10-Symptom-Rating"(ISR). [Development of the Questionnaire "ICD 10 Symptom Rating"(ISR)]. *Zeitschrift für Psychosomatische Medizin Und Psychotherapie [Journal for Psychosomatic Medicine and Psychotherapy]*, 54, 409–418. <https://doi.org/10.13109/zptm.2008.54.4.409>
- Tucker, L. R., & Lewis, C. (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika*, 38, 1–10. <https://doi.org/10.1007/BF02291170>
- Tzur-Bitan, D., Meiran, N., & Shahar, G. (2010). The importance of modeling comorbidity using an intra-individual, time-series approach. *Behavioral and Brain Sciences*, 33(2–3), 172–173. <https://doi.org/10.1017/s0140525x10000816>
- Tzur-Bitan, D., Meiran, N., Steinberg, D. M., & Shahar, G. (2012). Is the looming maladaptive cognitive style a central mechanism in the (Generalized) Anxiety-(Major) Depression comorbidity: An intra-individual time series study. *International Journal of Cognitive Therapy*, 5, 170–185. <https://doi.org/10.1521/ijct.2012.5.2.170>
- Wampold, B. (2015). Routine outcome monitoring: Coming of age with the usual developmental challenges. *Psychotherapy*, 52, 458–462. <https://doi.org/10.1037/pst0000037>
- Wampold, B. E., Flückiger, C., del Re, A. C., Yulish, N. E., Frost, N. D., Pace, B. T., ... Hilsenroth, M. J. (2017). In pursuit of truth: A critical examination of meta-analyses of cognitive behavior therapy. *Psychotherapy Research*, 27, 14–32. <https://doi.org/10.1080/10503307.2016.1249433>
- Wampold, B. E., & Imel, Z. E. (2015). *The great psychotherapy debate: The evidence for what makes psychotherapy work* (2nd ed.). New York: Routledge.
- Ware, J. E., Snow, K. K., Kosinski, M., & Gandek, B. (1993). *SF-36 Health Survey: Manual and interpretation guide*. Boston: The Health Institute, New England Medical Centre.
- Webber, C. L., & Zbilut, J. P. (1994). Dynamical assessment of physiological systems and states using recurrence plot strategies. *Journal of Applied Physiology*, 76, 965–973. <https://doi.org/10.1152/jap.1994.76.2.965>
- Wing, J., Beevor, A. S., Curtis, R. H., Park, S. B. G., Hadden, S., & Burns, A. (1998). Health of the Nation Outcome Scales (HoNOS): Research and development. *British Journal of Psychiatry*, 172, 11–18. <https://doi.org/10.1192/bjp.172.1.11>

How to cite this article: Schiepek G, Stöger-Schmidinger B, Kronberger H, et al. The Therapy Process Questionnaire - factor analysis and psychometric properties of a multidimensional self-rating scale for high-frequency monitoring of psychotherapeutic processes. *Clin Psychol Psychother*. 2019;1–17. <https://doi.org/10.1002/cpp.2384>